

3.4 Critical Pairs

Showing local confluence (Sketch):

Problem: If $t_1 \leftarrow_E t_0 \rightarrow_E t_2$, does there exist a term s such that $t_1 \rightarrow_E^* s \leftarrow_E^* t_2$?

If the two rewrite steps happen in different subtrees (disjoint redexes): yes.

If the two rewrite steps happen below each other (overlap at or below a variable position): yes.

If the left-hand sides of the two rules overlap at a non-variable position: needs further investigation.

Question:

Are there rewrite rules $l_1 \rightarrow r_1$ and $l_2 \rightarrow r_2$ such that some subterm l_1/p and l_2 have a common instance $(l_1/p)\sigma_1 = l_2\sigma_2$?

Observation:

If we assume w.o.l.o.g. that the two rewrite rules do not have common variables, then only a single substitution is necessary: $(l_1/p)\sigma = l_2\sigma$.

Further observation:

The mgu of l_1/p and l_2 subsumes all unifiers σ of l_1/p and l_2 .

Let $l_i \rightarrow r_i$ ($i = 1, 2$) be two rewrite rules in a TRS R whose variables have been renamed such that $\text{var}(l_1) \cap \text{var}(l_2) = \emptyset$. (Remember that $\text{var}(l_i) \supseteq \text{var}(r_i)$.)

Let $p \in \text{pos}(l_1)$ be a position such that l_1/p is not a variable and σ is an mgu of l_1/p and l_2 .

Then $r_1\sigma \leftarrow l_1\sigma \rightarrow (l_1\sigma)[r_2\sigma]_p$.

$\langle r_1\sigma, (l_1\sigma)[r_2\sigma]_p \rangle$ is called a *critical pair* of R .

The critical pair is *joinable* (or: converges), if $r_1\sigma \downarrow_R (l_1\sigma)[r_2\sigma]_p$.

Theorem 3.18 (“Critical Pair Theorem”) *A TRS R is locally confluent if and only if all its critical pairs are joinable.*

Proof. “only if”: obvious, since joinability of a critical pair is a special case of local confluence.

“if”: Suppose s rewrites to t_1 and t_2 using rewrite rules $l_i \rightarrow r_i \in R$ at positions $p_i \in \text{pos}(s)$, where $i = 1, 2$. Without loss of generality, we can assume that the two rules are variable disjoint, hence $s/p_i = l_i\theta$ and $t_i = s[r_i\theta]_{p_i}$.

We distinguish between two cases: Either p_1 and p_2 are in disjoint subtrees ($p_1 \parallel p_2$), or one is a prefix of the other (w.o.l.o.g., $p_1 \leq p_2$).

Case 1: $p_1 \parallel p_2$.

Then $s = s[l_1\theta]_{p_1}[l_2\theta]_{p_2}$, and therefore $t_1 = s[r_1\theta]_{p_1}[l_2\theta]_{p_2}$ and $t_2 = s[l_1\theta]_{p_1}[r_2\theta]_{p_2}$.

Let $t_0 = s[r_1\theta]_{p_1}[r_2\theta]_{p_2}$. Then clearly $t_1 \rightarrow_R t_0$ using $l_2 \rightarrow r_2$ and $t_2 \rightarrow_R t_0$ using $l_1 \rightarrow r_1$.

Case 2: $p_1 \leq p_2$.

Case 2.1: $p_2 = p_1q_1q_2$, where l_1/q_1 is some variable x .

In other words, the second rewrite step takes place at or below a variable in the first rule. Suppose that x occurs m times in l_1 and n times in r_1 (where $m \geq 1$ and $n \geq 0$).

Then $t_1 \rightarrow_R^* t_0$ by applying $l_2 \rightarrow r_2$ at all positions $p_1q'q_2$, where q' is a position of x in r_1 .

Conversely, $t_2 \rightarrow_R^* t_0$ by applying $l_2 \rightarrow r_2$ at all positions p_1qq_2 , where q is a position of x in l_1 different from q_1 , and by applying $l_1 \rightarrow r_1$ at p_1 with the substitution θ' , where $\theta' = \theta[x \mapsto (x\theta)[r_2\theta]_{q_2}]$.

Case 2.2: $p_2 = p_1p$, where p is a non-variable position of l_1 .

Then $s/p_2 = l_2\theta$ and $s/p_2 = (s/p_1)/p = (l_1\theta)/p = (l_1/p)\theta$, so θ is a unifier of l_2 and l_1/p .

Let σ be the mgu of l_2 and l_1/p , then $\theta = \tau \circ \sigma$ and $\langle r_1\sigma, (l_1\sigma)[r_2\sigma]_p \rangle$ is a critical pair.

By assumption, it is joinable, so $r_1\sigma \rightarrow_R^* v \leftarrow_R^* (l_1\sigma)[r_2\sigma]_p$.

Consequently, $t_1 = s[r_1\theta]_{p_1} = s[r_1\sigma\tau]_{p_1} \rightarrow_R^* s[v\tau]_{p_1}$ and $t_2 = s[r_2\theta]_{p_2} = s[(l_1\theta)[r_2\theta]_p]_{p_1} = s[(l_1\sigma\tau)[r_2\sigma\tau]_p]_{p_1} = s[((l_1\sigma)[r_2\sigma]_p)\tau]_{p_1} \rightarrow_R^* s[v\tau]_{p_1}$.

This completes the proof of the Critical Pair Theorem. □

Note: Critical pairs between a rule and (a renamed variant of) itself must be considered – except if the overlap is at the root (i. e., $p = \varepsilon$).

Corollary 3.19 *A terminating TRS R is confluent if and only if all its critical pairs are joinable.*

Proof. By Newman's Lemma and the Critical Pair Theorem. □

Corollary 3.20 *For a finite terminating TRS, confluence is decidable.*

Proof. For every pair of rules and every non-variable position in the first rule there is at most one critical pair $\langle u_1, u_2 \rangle$.

Reduce every u_i to some normal form u'_i . If $u'_1 = u'_2$ for every critical pair, then R is confluent, otherwise there is some non-confluent situation $u'_1 \leftarrow_R^* u_1 \leftarrow_R s \rightarrow_R u_2 \rightarrow_R^* u'_2$. □

3.5 Termination

Termination problems:

Given a finite TRS R and a term t , are all R -reductions starting from t terminating?

Given a finite TRS R , are all R -reductions terminating?

Proposition 3.21 *Both termination problems for TRSs are undecidable in general.*

Proof. Encode Turing machines using rewrite rules and reduce the (uniform) halting problems for TMs to the termination problems for TRSs. \square

Consequence:

Decidable criteria for termination are not complete.

Reduction Orderings

Goal:

Given a finite TRS R , show termination of R by looking at finitely many rules $l \rightarrow r \in R$, rather than at infinitely many possible replacement steps $s \rightarrow_R s'$.

A binary relation \sqsupset over $T_\Sigma(X)$ is called *compatible with Σ -operations*, if $s \sqsupset s'$ implies $f(t_1, \dots, s, \dots, t_n) \sqsupset f(t_1, \dots, s', \dots, t_n)$ for all $f \in \Omega$ and $s, s', t_i \in T_\Sigma(X)$.

Lemma 3.22 *The relation \sqsupset is compatible with Σ -operations, if and only if $s \sqsupset s'$ implies $t[s]_p \sqsupset t[s']_p$ for all $s, s', t \in T_\Sigma(X)$ and $p \in \text{pos}(t)$.*

Note: *compatible with Σ -operations* = *compatible with contexts*.

A binary relation \sqsupset over $T_\Sigma(X)$ is called *stable under substitutions*, if $s \sqsupset s'$ implies $s\sigma \sqsupset s'\sigma$ for all $s, s' \in T_\Sigma(X)$ and substitutions σ .

A binary relation \sqsupset is called a *rewrite relation*, if it is compatible with Σ -operations and stable under substitutions.

Example: If R is a TRS, then \rightarrow_R is a rewrite relation.

A strict partial ordering over $T_\Sigma(X)$ that is a rewrite relation is called *rewrite ordering*.

A well-founded rewrite ordering is called *reduction ordering*.

Theorem 3.23 *A TRS R terminates if and only if there exists a reduction ordering \succ such that $l \succ r$ for every rule $l \rightarrow r \in R$.*

Proof. “if”: $s \rightarrow_R s'$ if and only if $s = t[l\sigma]_p$, $s' = t[r\sigma]_p$. If $l \succ r$, then $l\sigma \succ r\sigma$ and therefore $t[l\sigma]_p \succ t[r\sigma]_p$. This implies $\rightarrow_R \subseteq \succ$. Since \succ is a well-founded ordering, \rightarrow_R is terminating.

“only if”: Define $\succ = \rightarrow_R^+$. If \rightarrow_R is terminating, then \succ is a reduction ordering. \square

The Interpretation Method

Proving termination by interpretation:

Let \mathcal{A} be a Σ -algebra; let \succ be a well-founded strict partial ordering on its universe.

Define the ordering $\succ_{\mathcal{A}}$ over $T_{\Sigma}(X)$ by $s \succ_{\mathcal{A}} t$ iff $\mathcal{A}(\beta)(s) \succ \mathcal{A}(\beta)(t)$ for all assignments $\beta : X \rightarrow U_{\mathcal{A}}$.

Is $\succ_{\mathcal{A}}$ a reduction ordering?

Lemma 3.24 $\succ_{\mathcal{A}}$ is stable under substitutions.

Proof. Let $s \succ_{\mathcal{A}} s'$, that is, $\mathcal{A}(\beta)(s) \succ \mathcal{A}(\beta)(s')$ for all assignments $\beta : X \rightarrow U_{\mathcal{A}}$. Let σ be a substitution. We have to show that $\mathcal{A}(\gamma)(s\sigma) \succ \mathcal{A}(\gamma)(s'\sigma)$ for all assignments $\gamma : X \rightarrow U_{\mathcal{A}}$. Choose $\beta = \gamma \circ \sigma$, then by the substitution lemma, $\mathcal{A}(\gamma)(s\sigma) = \mathcal{A}(\beta)(s) \succ \mathcal{A}(\beta)(s') = \mathcal{A}(\gamma)(s'\sigma)$. Therefore $s\sigma \succ_{\mathcal{A}} s'\sigma$. \square

A function $F : U_{\mathcal{A}}^n \rightarrow U_{\mathcal{A}}$ is called *monotone* (with respect to \succ), if $a \succ a'$ implies $F(b_1, \dots, a, \dots, b_n) \succ F(b_1, \dots, a', \dots, b_n)$ for all $a, a', b_i \in U_{\mathcal{A}}$.

Lemma 3.25 If the interpretation $f_{\mathcal{A}}$ of every function symbol f is monotone w. r. t. \succ , then $\succ_{\mathcal{A}}$ is compatible with Σ -operations.

Proof. Let $s \succ s'$, that is, $\mathcal{A}(\beta)(s) \succ \mathcal{A}(\beta)(s')$ for all $\beta : X \rightarrow U_{\mathcal{A}}$. Let $\beta : X \rightarrow U_{\mathcal{A}}$ be an arbitrary assignment. Then

$$\begin{aligned} \mathcal{A}(\beta)(f(t_1, \dots, s, \dots, t_n)) &= f_{\mathcal{A}}(\mathcal{A}(\beta)(t_1), \dots, \mathcal{A}(\beta)(s), \dots, \mathcal{A}(\beta)(t_n)) \\ &\succ f_{\mathcal{A}}(\mathcal{A}(\beta)(t_1), \dots, \mathcal{A}(\beta)(s'), \dots, \mathcal{A}(\beta)(t_n)) \\ &= \mathcal{A}(\beta)(f(t_1, \dots, s', \dots, t_n)) \end{aligned}$$

Therefore $f(t_1, \dots, s, \dots, t_n) \succ_{\mathcal{A}} f(t_1, \dots, s', \dots, t_n)$. \square

Theorem 3.26 If the interpretation $f_{\mathcal{A}}$ of every function symbol f is monotone w. r. t. \succ , then $\succ_{\mathcal{A}}$ is a reduction ordering.

Proof. By the previous two lemmas, $\succ_{\mathcal{A}}$ is a rewrite relation. If there were an infinite chain $s_1 \succ_{\mathcal{A}} s_2 \succ_{\mathcal{A}} \dots$, then it would correspond to an infinite chain $\mathcal{A}(\beta)(s_1) \succ \mathcal{A}(\beta)(s_2) \succ \dots$ (with β chosen arbitrarily). Thus $\succ_{\mathcal{A}}$ is well-founded. Irreflexivity and transitivity are proved similarly. \square

Polynomial Orderings

Polynomial orderings:

Instance of the interpretation method:

The carrier set $U_{\mathcal{A}}$ is some subset of the natural numbers.

To every function symbol f with arity n we associate a polynomial $P_f(X_1, \dots, X_n) \in \mathbb{N}[X_1, \dots, X_n]$ with coefficients in \mathbb{N} and indeterminates X_1, \dots, X_n . Then we define $f_{\mathcal{A}}(a_1, \dots, a_n) = P_f(a_1, \dots, a_n)$ for $a_i \in U_{\mathcal{A}}$.

Requirement 1:

If $a_1, \dots, a_n \in U_{\mathcal{A}}$, then $f_{\mathcal{A}}(a_1, \dots, a_n) \in U_{\mathcal{A}}$. (Otherwise, \mathcal{A} would not be a Σ -algebra.)

Requirement 2:

$f_{\mathcal{A}}$ must be monotone (w.r.t. \succ).

From now on:

$$U_{\mathcal{A}} = \{n \in \mathbb{N} \mid n \geq 2\}.$$

If $\text{arity}(f) = 0$, then P_f is a constant ≥ 2 .

If $\text{arity}(f) = n \geq 1$, then P_f is a polynomial $P(X_1, \dots, X_n)$, such that every X_i occurs in some monomial with exponent at least 1 and non-zero coefficient.

\Rightarrow Requirements 1 and 2 are satisfied.

The mapping from function symbols to polynomials can be extended to terms: A term t containing the variables x_1, \dots, x_n yields a polynomial P_t with indeterminates X_1, \dots, X_n (where X_i corresponds to $\beta(x_i)$).

Example:

$$\Omega = \{b, f, g\} \text{ with } \text{arity}(b) = 0, \text{arity}(f) = 1, \text{arity}(g) = 3,$$

$$U_{\mathcal{A}} = \{n \in \mathbb{N} \mid n \geq 2\},$$

$$P_b = 3, \quad P_f(X_1) = X_1^2, \quad P_g(X_1, X_2, X_3) = X_1 + X_2X_3.$$

$$\text{Let } t = g(f(b), f(x), y), \text{ then } P_t(X, Y) = 9 + X^2Y.$$

If P, Q are polynomials in $\mathbb{N}[X_1, \dots, X_n]$, we write $P > Q$ if $P(a_1, \dots, a_n) > Q(a_1, \dots, a_n)$ for all $a_1, \dots, a_n \in U_{\mathcal{A}}$.

Clearly, $l \succ_{\mathcal{A}} r$ iff $P_l > P_r$.

Question: Can we check $P_l > P_r$ automatically?

Hilbert's 10th Problem:

Given a polynomial $P \in \mathbb{Z}[X_1, \dots, X_n]$ with integer coefficients, is $P = 0$ for some n -tuple of natural numbers?

Theorem 3.27 *Hilbert's 10th Problem is undecidable.*

Proposition 3.28 *Given a polynomial interpretation and two terms l, r , it is undecidable whether $P_l > P_r$.*

Proof. By reduction of Hilbert's 10th Problem. □

One possible solution:

Test whether $P_l(a_1, \dots, a_n) > P_r(a_1, \dots, a_n)$ for all $a_1, \dots, a_n \in \{x \in \mathbb{R} \mid x \geq 2\}$.

This is decidable (but very slow). Since $U_{\mathcal{A}} \subseteq \{x \in \mathbb{R} \mid x \geq 2\}$, it implies $P_l > P_r$.

Another solution (Ben Cherifa and Lescanne):

Consider the difference $P_l(X_1, \dots, X_n) - P_r(X_1, \dots, X_n)$ as a polynomial with real coefficients and apply the following inference system to it to show that it is positive for all $a_1, \dots, a_n \in U_{\mathcal{A}}$:

$$P \Rightarrow_{BCL} \top,$$

if P contains at least one monomial with a positive coefficient and no monomial with a negative coefficient.

$$P + cX_1^{p_1} \dots X_n^{p_n} - dX_1^{q_1} \dots X_n^{q_n} \Rightarrow_{BCL} P + c'X_1^{p_1} \dots X_n^{p_n},$$

if $c, d > 0$, $p_i \geq q_i$ for all i , and $c' = c - d \cdot 2^{(q_1 - p_1) + \dots + (q_n - p_n)} \geq 0$.

$$P + cX_1^{p_1} \dots X_n^{p_n} - dX_1^{q_1} \dots X_n^{q_n} \Rightarrow_{BCL} P - d'X_1^{q_1} \dots X_n^{q_n},$$

if $c, d > 0$, $p_i \geq q_i$ for all i , and $d' = d - c \cdot 2^{(p_1 - q_1) + \dots + (p_n - q_n)} > 0$.

Lemma 3.29 *If $P \Rightarrow_{BCL} P'$, then $P(a_1, \dots, a_n) \geq P'(a_1, \dots, a_n)$ for all $a_1, \dots, a_n \in U_{\mathcal{A}}$.*

Proof. Follows from the fact that $a_i \in U_{\mathcal{A}}$ implies $a_i \geq 2$. □

Proposition 3.30 *If $P \Rightarrow_{BCL}^+ \top$, then $P(a_1, \dots, a_n) > 0$ for all $a_1, \dots, a_n \in U_{\mathcal{A}}$.*

Simplification Orderings

The *proper subterm ordering* \triangleright is defined by $s \triangleright t$ if and only if $s/p = t$ for some position $p \neq \varepsilon$ of s .

A rewrite ordering \succ over $T_\Sigma(X)$ is called *simplification ordering*, if it has the *subterm property*: $s \triangleright t$ implies $s \succ t$ for all $s, t \in T_\Sigma(X)$.

Example:

Let R_{emb} be the rewrite system $R_{\text{emb}} = \{f(x_1, \dots, x_n) \rightarrow x_i \mid f \in \Omega, 1 \leq i \leq n = \text{arity}(f)\}$.

Define $\triangleright_{\text{emb}} = \rightarrow_{R_{\text{emb}}}^+$ and $\succeq_{\text{emb}} = \rightarrow_{R_{\text{emb}}}^*$ (“homeomorphic embedding relation”).

$\triangleright_{\text{emb}}$ is a simplification ordering.

Lemma 3.31 *If \succ is a simplification ordering, then $s \triangleright_{\text{emb}} t$ implies $s \succ t$ and $s \succeq_{\text{emb}} t$ implies $s \succeq t$.*

Proof. Since \succ is transitive and \succeq is transitive and reflexive, it suffices to show that $s \rightarrow_{R_{\text{emb}}} t$ implies $s \succ t$. By definition, $s \rightarrow_{R_{\text{emb}}} t$ if and only if $s = s[l\sigma]$ and $t = s[r\sigma]$ for some rule $l \rightarrow r \in R_{\text{emb}}$. Obviously, $l \triangleright r$ for all rules in R_{emb} , hence $l \succ r$. Since \succ is a rewrite relation, $s = s[l\sigma] \succ s[r\sigma] = t$. \square

Goal:

Show that every simplification ordering is well-founded (and therefore a reduction ordering).

Note: This works only for *finite* signatures!

To fix this for infinite signatures, the definition of simplification orderings and the definition of embedding have to be modified.

Theorem 3.32 (“Kruskal’s Theorem”) *Let Σ be a finite signature, let X be a finite set of variables. Then for every infinite sequence t_1, t_2, t_3, \dots there are indices $j > i$ such that $t_j \succeq_{\text{emb}} t_i$. (\succeq_{emb} is called a well-partial-ordering (wpo).)*

Proof. See Baader and Nipkow, page 113–115. \square

Theorem 3.33 (Dershowitz) *If Σ is a finite signature, then every simplification ordering \succ on $T_\Sigma(X)$ is well-founded (and therefore a reduction ordering).*

Proof. Suppose that $t_1 \succ t_2 \succ t_3 \succ \dots$ is an infinite descending chain.

First assume that there is an $x \in \text{var}(t_{i+1}) \setminus \text{var}(t_i)$. Let $\sigma = [t_i/x]$, then $t_{i+1}\sigma \supseteq x\sigma = t_i$ and therefore $t_i = t_i\sigma \succ t_{i+1}\sigma \succeq t_i$, contradicting reflexivity.

Consequently, $\text{var}(t_i) \supseteq \text{var}(t_{i+1})$ and $t_i \in T_\Sigma(V)$ for all i , where V is the finite set $\text{var}(t_1)$. By Kruskal's Theorem, there are $i < j$ with $t_i \trianglelefteq_{\text{emb}} t_j$. Hence $t_i \preceq t_j$, contradicting $t_i \succ t_j$. \square

There are reduction orderings that are not simplification orderings and terminating TRSs that are not contained in any simplification ordering.

Example:

Let $R = \{f(f(x)) \rightarrow f(g(f(x)))\}$.

R terminates and \rightarrow_R^+ is therefore a reduction ordering.

Assume that \rightarrow_R were contained in a simplification ordering \succ . Then $f(f(x)) \rightarrow_R f(g(f(x)))$ implies $f(f(x)) \succ f(g(f(x)))$, and $f(g(f(x))) \supseteq_{\text{emb}} f(f(x))$ implies $f(g(f(x))) \succeq f(f(x))$, hence $f(f(x)) \succ f(f(x))$.

Recursive Path Orderings

Let $\Sigma = (\Omega, \Pi)$ be a finite signature, let \succ be a strict partial ordering (“precedence”) on Ω .

The *lexicographic path ordering* \succ_{lpo} on $T_\Sigma(X)$ induced by \succ is defined by: $s \succ_{\text{lpo}} t$ iff

- (1) $t \in \text{var}(s)$ and $t \neq s$, or
- (2) $s = f(s_1, \dots, s_m)$, $t = g(t_1, \dots, t_n)$, and
 - (a) $s_i \succeq_{\text{lpo}} t$ for some i , or
 - (b) $f \succ g$ and $s \succ_{\text{lpo}} t_j$ for all j , or
 - (c) $f = g$, $s \succ_{\text{lpo}} t_j$ for all j , and $(s_1, \dots, s_m) (\succ_{\text{lpo}})_{\text{lex}} (t_1, \dots, t_n)$.

Lemma 3.34 *$s \succ_{\text{lpo}} t$ implies $\text{var}(s) \supseteq \text{var}(t)$.*

Proof. By induction on $|s| + |t|$ and case analysis. \square

Theorem 3.35 \succ_{lpo} is a simplification ordering on $T_{\Sigma}(X)$.

Proof. Show transitivity, subterm property, stability under substitutions, compatibility with Σ -operations, and irreflexivity, usually by induction on the sum of the term sizes and case analysis. Details: Baader and Nipkow, page 119/120. \square

Theorem 3.36 If the precedence \succ is total, then the lexicographic path ordering \succ_{lpo} is total on ground terms, i. e., for all $s, t \in T_{\Sigma}(\emptyset)$: $s \succ_{\text{lpo}} t \vee t \succ_{\text{lpo}} s \vee s = t$.

Proof. By induction on $|s| + |t|$ and case analysis. \square

Recapitulation:

Let $\Sigma = (\Omega, \Pi)$ be a finite signature, let \succ be a strict partial ordering (“precedence”) on Ω . The *lexicographic path ordering* \succ_{lpo} on $T_{\Sigma}(X)$ induced by \succ is defined by: $s \succ_{\text{lpo}} t$ iff

- (1) $t \in \text{var}(s)$ and $t \neq s$, or
- (2) $s = f(s_1, \dots, s_m)$, $t = g(t_1, \dots, t_n)$, and
 - (a) $s_i \succeq_{\text{lpo}} t$ for some i , or
 - (b) $f \succ g$ and $s \succ_{\text{lpo}} t_j$ for all j , or
 - (c) $f = g$, $s \succ_{\text{lpo}} t_j$ for all j , and $(s_1, \dots, s_m) (\succ_{\text{lpo}})_{\text{lex}} (t_1, \dots, t_n)$.

There are several possibilities to compare subterms in (2)(c):

compare list of subterms lexicographically left-to-right (“*lexicographic path ordering (lpo)*”, Kamin and Lévy)

compare list of subterms lexicographically right-to-left (or according to some permutation π)

compare multiset of subterms using the multiset extension (“*multiset path ordering (mpo)*”, Dershowitz)

to each function symbol f with $\text{arity}(n) \geq 1$ associate a status $\in \{\text{mul}\} \cup \{\text{lex}_{\pi} \mid \pi : \{1, \dots, n\} \rightarrow \{1, \dots, n\}\}$ and compare according to that status (“*recursive path ordering (rpo) with status*”)

The Knuth-Bendix Ordering

Let $\Sigma = (\Omega, \Pi)$ be a finite signature, let \succ be a strict partial ordering (“precedence”) on Ω , let $w : \Omega \cup X \rightarrow \mathbb{R}_0^+$ be a *weight function*, such that the following admissibility conditions are satisfied:

$w(x) = w_0 \in \mathbb{R}^+$ for all variables $x \in X$; $w(c) \geq w_0$ for all constants $c \in \Omega$.

If $w(f) = 0$ for some $f \in \Omega$ with $\text{arity}(f) = 1$, then $f \succeq g$ for all $g \in \Omega$.

The weight function w can be extended to terms as follows:

$$w(t) = \sum_{x \in \text{var}(t)} w(x) \cdot \#(x, t) + \sum_{f \in \Omega} w(f) \cdot \#(f, t).$$

The *Knuth-Bendix ordering* \succ_{kbo} on $T_\Sigma(X)$ induced by \succ and w is defined by: $s \succ_{\text{kbo}} t$ iff

- (1) $\#(x, s) \geq \#(x, t)$ for all variables x and $w(s) > w(t)$, or
- (2) $\#(x, s) \geq \#(x, t)$ for all variables x , $w(s) = w(t)$, and
 - (a) $t = x$, $s = f^n(x)$ for some $n \geq 1$, or
 - (b) $s = f(s_1, \dots, s_m)$, $t = g(t_1, \dots, t_n)$, and $f \succ g$, or
 - (c) $s = f(s_1, \dots, s_m)$, $t = f(t_1, \dots, t_m)$, and $(s_1, \dots, s_m) (\succ_{\text{kbo}})_{\text{lex}} (t_1, \dots, t_m)$.

Theorem 3.37 *The Knuth-Bendix ordering induced by \succ and w is a simplification ordering on $T_\Sigma(X)$.*

Proof. Baader and Nipkow, pages 125–129. □