

Case 3.2: $s\theta \succ s'\theta$.

If $s\theta \downarrow_{R_{C\theta}} s'\theta$ and $s\theta \succ s'\theta$, then $s\theta$ must be reducible by some rule in some $E_{D\theta} \subseteq R_{C\theta}$. (Without loss of generality we assume that C and D are variable disjoint; so we can use the same substitution θ .) Let $D\theta = D'\theta \vee t\theta \approx t'\theta$ with $E_{D\theta} = \{t\theta \rightarrow t'\theta\}$. Since $D\theta$ is productive, $D'\theta$ is false in $R_{C\theta}$. Besides, by part (ii) of the induction hypothesis, $D\theta$ is not redundant w.r.t. $G_\Sigma(N)$, so D is not redundant w.r.t. N . Note that $t\theta$ cannot occur in $s\theta$ at or below a variable position of s , say $x\theta = w[t\theta]$, since otherwise $C\theta$ would be subject to Case 2 above. Consequently, the *left superposition* inference

$$\frac{D'\theta \vee t\theta \approx t'\theta \quad C'\theta \vee s\theta[t\theta] \not\approx s'\theta}{D'\theta \vee C'\theta \vee s\theta[t'\theta] \not\approx s'\theta}$$

is a ground instance of a *left superposition* inference from D and C . By saturation up to redundancy, its conclusion is either contained in $G_\Sigma(N)$ and smaller than $C\theta$, or it follows from clauses in $G_\Sigma(N)$ that are smaller than itself (and therefore smaller than $C\theta$). By the induction hypothesis, these clauses are true in $R_{C\theta}$, thus $D'\theta \vee C'\theta \vee s\theta[t'\theta] \not\approx s'\theta$ is true in $R_{C\theta}$. Since $D'\theta$ and $s\theta[t'\theta] \not\approx s'\theta$ are false in $R_{C\theta}$, both $C'\theta$ and $C\theta$ must be true.

Case 4: $C\theta$ does not contain a maximal negative literal.

Suppose that $C\theta$ does not fall into Cases 1 to 3. Then $C\theta$ can be written as $C'\theta \vee s\theta \approx s'\theta$, where $s\theta \approx s'\theta$ is a maximal literal of $C\theta$. If $E_{C\theta} = \{s\theta \rightarrow s'\theta\}$ or $C'\theta$ is true in $R_{C\theta}$ or $s\theta = s'\theta$, then there is nothing to show, so assume that $E_{C\theta} = \emptyset$ and that $C'\theta$ is false in $R_{C\theta}$. Without loss of generality, $s\theta \succ s'\theta$.

Case 4.1: $s\theta \approx s'\theta$ is maximal in $C\theta$, but not strictly maximal.

If $s\theta \approx s'\theta$ is maximal in $C\theta$, but not strictly maximal, then $C\theta$ can be written as $C''\theta \vee t\theta \approx t'\theta \vee s\theta \approx s'\theta$, where $t\theta = s\theta$ and $t'\theta = s'\theta$. In this case, there is a *equality factoring* inference

$$\frac{C''\theta \vee t\theta \approx t'\theta \vee s\theta \approx s'\theta}{C''\theta \vee t'\theta \not\approx s'\theta \vee t\theta \approx t'\theta}$$

This inference is a ground instance of an inference from C . By induction hypothesis, its conclusion is true in $R_{C\theta}$. Trivially, $t'\theta = s'\theta$ implies $t'\theta \downarrow_{R_{C\theta}} s'\theta$, so $t'\theta \not\approx s'\theta$ must be false and $C\theta$ must be true in $R_{C\theta}$.

Case 4.2: $s\theta \approx s'\theta$ is strictly maximal in $C\theta$ and $s\theta$ is reducible.

Suppose that $s\theta \approx s'\theta$ is strictly maximal in $C\theta$ and $s\theta$ is reducible by some rule in $E_{D\theta} \subseteq R_{C\theta}$. Let $D\theta = D'\theta \vee t\theta \approx t'\theta$ and $E_{D\theta} = \{t\theta \rightarrow t'\theta\}$. Since $D\theta$ is productive, $D\theta$ is not redundant and $D'\theta$ is false in $R_{C\theta}$. We can now proceed in essentially the

same way as in Case 3.2: If $t\theta$ occurred in $s\theta$ at or below a variable position of s , say $x\theta = w[t\theta]$, then $C\theta$ would be subject to Case 2 above. Otherwise, the *right superposition* inference

$$\frac{D'\theta \vee t\theta \approx t'\theta \quad C'\theta \vee s\theta[t\theta] \approx s'\theta}{D'\theta \vee C'\theta \vee s\theta[t'\theta] \approx s'\theta}$$

is a ground instance of a *right superposition* inference from D and C . By saturation up to redundancy, its conclusion is true in $R_{C\theta}$. Since $D'\theta$ and $C'\theta$ are false in $R_{C\theta}$, $s\theta[t'\theta] \approx s'\theta$ must be true in $R_{C\theta}$. On the other hand, $t\theta \approx t'\theta$ is true in $R_{C\theta}$, so by congruence, $s\theta[t\theta] \approx s'\theta$ and $C\theta$ are true in $R_{C\theta}$.

Case 4.3: $s\theta \approx s'\theta$ is strictly maximal in $C\theta$ and $s\theta$ is irreducible.

Suppose that $s\theta \approx s'\theta$ is strictly maximal in $C\theta$ and $s\theta$ is irreducible by $R_{C\theta}$. Then there are three possibilities: $C\theta$ can be true in $R_{C\theta}$, or $C'\theta$ can be true in $R_{C\theta} \cup \{s\theta \rightarrow s'\theta\}$, or $E_{C\theta} = \{s\theta \rightarrow s'\theta\}$. In the first and the third case, there is nothing to show. Let us therefore assume that $C\theta$ is false in $R_{C\theta}$ and $C'\theta$ is true in $R_{C\theta} \cup \{s\theta \rightarrow s'\theta\}$. Then $C'\theta = C''\theta \vee t\theta \approx t'\theta$, where the literal $t\theta \approx t'\theta$ is true in $R_{C\theta} \cup \{s\theta \rightarrow s'\theta\}$ and false in $R_{C\theta}$. In other words, $t\theta \downarrow_{R_{C\theta} \cup \{s\theta \rightarrow s'\theta\}} t'\theta$, but not $t\theta \downarrow_{R_{C\theta}} t'\theta$. Consequently, there is a rewrite proof of $t\theta \rightarrow^* u \leftarrow^* t'\theta$ by $R_{C\theta} \cup \{s\theta \rightarrow s'\theta\}$ in which the rule $s\theta \rightarrow s'\theta$ is used at least once. Without loss of generality we assume that $t\theta \succeq t'\theta$. Since $s\theta \approx s'\theta \succ_L t\theta \approx t'\theta$ and $s\theta \succ s'\theta$ we can conclude that $s\theta \succeq t\theta \succ t'\theta$. But then there is only one possibility how the rule $s\theta \rightarrow s'\theta$ can be used in the rewrite proof: We must have $s\theta = t\theta$ and the rewrite proof must have the form $t\theta \rightarrow s'\theta \rightarrow^* u \leftarrow^* t'\theta$, where the first step uses $s\theta \rightarrow s'\theta$ and all other steps use rules from $R_{C\theta}$. Consequently, $s'\theta \approx t'\theta$ is true in $R_{C\theta}$. Now observe that there is an *equality factoring* inference

$$\frac{C''\theta \vee t\theta \approx t'\theta \vee s\theta \approx s'\theta}{C''\theta \vee t'\theta \not\approx s'\theta \vee t\theta \approx t'\theta}$$

whose conclusion is true in $R_{C\theta}$ by saturation. Since the literal $t'\theta \not\approx s'\theta$ must be false in $R_{C\theta}$, the rest of the clause must be true in $R_{C\theta}$, and therefore $C\theta$ must be true in $R_{C\theta}$, contradicting our assumption. This concludes the proof of the theorem. $\square$

A Σ -interpretation $\mathcal{A}$ is called *term-generated*, if for every $b \in U_{\mathcal{A}}$ there is a ground term $t \in T_{\Sigma}(\emptyset)$ such that $b = \mathcal{A}(\beta)(t)$.

Lemma 4.53 *Let N be a set of (universally quantified) Σ -clauses and let $\mathcal{A}$ be a term-generated Σ -interpretation. Then $\mathcal{A}$ is a model of $G_{\Sigma}(N)$ if and only if it is a model of N .*

Proof. ($\Rightarrow$): Let $\mathcal{A} \models G_\Sigma(N)$; let $(\forall \vec{x}C) \in N$. Then $\mathcal{A} \models \forall \vec{x}C$ iff $\mathcal{A}(\gamma[x_i \mapsto a_i])(C) = 1$ for all γ and a_i . Choose ground terms t_i such that $\mathcal{A}(\gamma)(t_i) = a_i$; define θ such that $x_i\theta = t_i$, then $\mathcal{A}(\gamma[x_i \mapsto a_i])(C) = \mathcal{A}(\gamma \circ \theta)(C) = \mathcal{A}(\gamma)(C\theta) = 1$ since $C\theta \in G_\Sigma(N)$.

($\Leftarrow$): Let $\mathcal{A}$ be a model of N ; let $C \in N$ and $C\theta \in G_\Sigma(N)$. Then $\mathcal{A}(\gamma)(C\theta) = \mathcal{A}(\gamma \circ \theta)(C) = 1$ since $\mathcal{A} \models N$. $\square$

Theorem 4.54 (Refutational Completeness: Static View) *Let N be a set of clauses that is saturated up to redundancy. Then N has a model if and only if N does not contain the empty clause.*

Proof. If $\perp \in N$, then obviously N does not have a model. If $\perp \notin N$, then the interpretation R_∞ (that is, $T_\Sigma(\emptyset)/R_\infty$) is a model of all ground instances in $G_\Sigma(N)$ according to part (iii) of the model construction theorem. As $T_\Sigma(\emptyset)/R_\infty$ is term generated, it is a model of N . $\square$

So far, we have considered only inference rules that add new clauses to the current set of clauses (corresponding to the *Deduce* rule of Knuth-Bendix Completion).

In other words, we have derivations of the form $N_0 \vdash N_1 \vdash N_2 \vdash \dots$, where each N_{i+1} is obtained from N_i by adding the consequence of some inference from clauses in N_i .

Under which circumstances are we allowed to delete (or simplify) a clause during the derivation?

A *run* of the superposition calculus is a sequence $N_0 \vdash N_1 \vdash N_2 \vdash \dots$, such that

- (i) $N_i \models N_{i+1}$, and
- (ii) all clauses in $N_i \setminus N_{i+1}$ are redundant w.r.t. N_{i+1} .

In other words, during a run we may add a new clause if it follows from the old ones, and we may delete a clause, if it is redundant w.r.t. the remaining ones.

For a run, $N_\infty = \bigcup_{i \geq 0} N_i$ and $N_* = \bigcup_{i \geq 0} \bigcap_{j \geq i} N_j$. The set N_* of all *persistent* clauses is called the *limit* of the run.

Lemma 4.55 *If $N \subseteq N'$, then $Red(N) \subseteq Red(N')$.*

Proof. Obvious. $\square$

Lemma 4.56 *If $N' \subseteq Red(N)$, then $Red(N) \subseteq Red(N \setminus N')$.*

Proof. Follows from the compactness of first-order logic and the well-foundedness of the multiset extension of the clause ordering. $\square$