

4.5 Termination

Termination problems:

Given a finite TRS R and a term t , are all R -reductions starting from t terminating?

Given a finite TRS R , are all R -reductions terminating?

Proposition 4.21 *Both termination problems for TRSs are undecidable in general.*

Proof. Encode Turing machines using rewrite rules and reduce the (uniform) halting problems for TMs to the termination problems for TRSs. \square

Consequence:

Decidable criteria for termination are not complete.

Reduction Orderings

Goal:

Given a finite TRS R , show termination of R by looking at finitely many rules $l \rightarrow r \in R$, rather than at infinitely many possible replacement steps $s \rightarrow_R s'$.

A binary relation \sqsupset over $T_\Sigma(X)$ is called *compatible with Σ -operations*, if $s \sqsupset s'$ implies $f(t_1, \dots, s, \dots, t_n) \sqsupset f(t_1, \dots, s', \dots, t_n)$ for all $f \in \Omega$ and $s, s', t_i \in T_\Sigma(X)$.

Lemma 4.22 *The relation \sqsupset is compatible with Σ -operations, if and only if $s \sqsupset s'$ implies $t[s]_p \sqsupset t[s']_p$ for all $s, s', t \in T_\Sigma(X)$ and $p \in \text{pos}(t)$.*

Note: *compatible with Σ -operations* = *compatible with contexts*.

A binary relation \sqsupset over $T_\Sigma(X)$ is called *stable under substitutions*, if $s \sqsupset s'$ implies $s\sigma \sqsupset s'\sigma$ for all $s, s' \in T_\Sigma(X)$ and substitutions σ .

A binary relation \sqsupset is called a *rewrite relation*, if it is compatible with Σ -operations and stable under substitutions.

Example: If R is a TRS, then \rightarrow_R is a rewrite relation.

A strict partial ordering over $T_\Sigma(X)$ that is a rewrite relation is called *rewrite ordering*.

A well-founded rewrite ordering is called *reduction ordering*.

Theorem 4.23 *A TRS R terminates if and only if there exists a reduction ordering \succ such that $l \succ r$ for every rule $l \rightarrow r \in R$.*

Proof. “if”: $s \rightarrow_R s'$ if and only if $s = t[l\sigma]_p$, $s' = t[r\sigma]_p$. If $l \succ r$, then $l\sigma \succ r\sigma$ and therefore $t[l\sigma]_p \succ t[r\sigma]_p$. This implies $\rightarrow_R \subseteq \succ$. Since \succ is a well-founded ordering, \rightarrow_R is terminating.

“only if”: Define $\succ = \rightarrow_R^+$. If \rightarrow_R is terminating, then \succ is a reduction ordering. \square

Simplification Orderings

The *proper subterm ordering* \triangleright is defined by $s \triangleright t$ if and only if $s/p = t$ for some position $p \neq \varepsilon$ of s .

A rewrite ordering \succ over $T_\Sigma(X)$ is called *simplification ordering*, if it has the *subterm property*: $s \triangleright t$ implies $s \succ t$ for all $s, t \in T_\Sigma(X)$.

Example:

Let R_{emb} be the rewrite system $R_{\text{emb}} = \{f(x_1, \dots, x_n) \rightarrow x_i \mid f \in \Omega, 1 \leq i \leq n = \text{arity}(f)\}$.

Define $\triangleright_{\text{emb}} = \rightarrow_{R_{\text{emb}}}^+$ and $\succeq_{\text{emb}} = \rightarrow_{R_{\text{emb}}}^*$ (“homeomorphic embedding relation”).

$\triangleright_{\text{emb}}$ is a simplification ordering.

Lemma 4.24 *If \succ is a simplification ordering, then $s \triangleright_{\text{emb}} t$ implies $s \succ t$ and $s \succeq_{\text{emb}} t$ implies $s \succeq t$.*

Proof. Since \succ is transitive and \succeq is transitive and reflexive, it suffices to show that $s \rightarrow_{R_{\text{emb}}} t$ implies $s \succ t$. By definition, $s \rightarrow_{R_{\text{emb}}} t$ if and only if $s = s[l\sigma]$ and $t = s[r\sigma]$ for some rule $l \rightarrow r \in R_{\text{emb}}$. Obviously, $l \triangleright r$ for all rules in R_{emb} , hence $l \succ r$. Since \succ is a rewrite relation, $s = s[l\sigma] \succ s[r\sigma] = t$. \square

Goal:

Show that every simplification ordering is well-founded (and therefore a reduction ordering).

Note: This works only for *finite* signatures!

To fix this for infinite signatures, the definition of simplification orderings and the definition of embedding have to be modified.

Theorem 4.25 (“Kruskal’s Theorem”) *Let Σ be a finite signature, let X be a finite set of variables. Then for every infinite sequence t_1, t_2, t_3, \dots there are indices $j > i$ such that $t_j \succeq_{\text{emb}} t_i$. (\succeq_{emb} is called a well-partial-ordering (wpo).)*

Proof. See Baader and Nipkow, page 113–115. \square

Theorem 4.26 (Dershowitz) *If Σ is a finite signature, then every simplification ordering \succ on $T_\Sigma(X)$ is well-founded (and therefore a reduction ordering).*

Proof. Suppose that $t_1 \succ t_2 \succ t_3 \succ \dots$ is an infinite descending chain.

First assume that there is an $x \in \text{var}(t_{i+1}) \setminus \text{var}(t_i)$. Let $\sigma = [t_i/x]$, then $t_{i+1}\sigma \supseteq x\sigma = t_i$ and therefore $t_i = t_i\sigma \succ t_{i+1}\sigma \succeq t_i$, contradicting reflexivity.

Consequently, $\text{var}(t_i) \supseteq \text{var}(t_{i+1})$ and $t_i \in T_\Sigma(V)$ for all i , where V is the finite set $\text{var}(t_1)$. By Kruskal's Theorem, there are $i < j$ with $t_i \preceq_{\text{emb}} t_j$. Hence $t_i \preceq t_j$, contradicting $t_i \succ t_j$. \square

There are reduction orderings that are not simplification orderings and terminating TRSs that are not contained in any simplification ordering.

Example:

Let $R = \{f(f(x)) \rightarrow f(g(f(x)))\}$.

R terminates and \rightarrow_R^+ is therefore a reduction ordering.

Assume that \rightarrow_R were contained in a simplification ordering \succ . Then $f(f(x)) \rightarrow_R f(g(f(x)))$ implies $f(f(x)) \succ f(g(f(x)))$, and $f(g(f(x))) \supseteq_{\text{emb}} f(f(x))$ implies $f(g(f(x))) \succeq f(f(x))$, hence $f(f(x)) \succ f(f(x))$.

Recursive Path Orderings

Let $\Sigma = (\Omega, \Pi)$ be a finite signature, let \succ be a strict partial ordering (“precedence”) on Ω .

The *lexicographic path ordering* \succ_{lpo} on $T_\Sigma(X)$ induced by \succ is defined by: $s \succ_{\text{lpo}} t$ iff

- (1) $t \in \text{var}(s)$ and $t \neq s$, or
- (2) $s = f(s_1, \dots, s_m)$, $t = g(t_1, \dots, t_n)$, and
 - (a) $s_i \succeq_{\text{lpo}} t$ for some i , or
 - (b) $f \succ g$ and $s \succ_{\text{lpo}} t_j$ for all j , or
 - (c) $f = g$, $s \succ_{\text{lpo}} t_j$ for all j , and $(s_1, \dots, s_m) (\succ_{\text{lpo}})_{\text{lex}} (t_1, \dots, t_n)$.

Lemma 4.27 $s \succ_{\text{lpo}} t$ implies $\text{var}(s) \supseteq \text{var}(t)$.

Proof. By induction on $|s| + |t|$ and case analysis. \square

Theorem 4.28 \succ_{lpo} is a simplification ordering on $T_\Sigma(X)$.

Proof. Show transitivity, subterm property, stability under substitutions, compatibility with Σ -operations, and irreflexivity, usually by induction on the sum of the term sizes and case analysis. Details: Baader and Nipkow, page 119/120. \square

Theorem 4.29 *If the precedence \succ is total, then the lexicographic path ordering \succ_{lpo} is total on ground terms, i. e., for all $s, t \in \mathsf{T}_{\Sigma}(\emptyset)$: $s \succ_{\text{lpo}} t \vee t \succ_{\text{lpo}} s \vee s = t$.*

Proof. By induction on $|s| + |t|$ and case analysis. \square

Recapitulation:

Let $\Sigma = (\Omega, \Pi)$ be a finite signature, let \succ be a strict partial ordering (“precedence”) on Ω . The *lexicographic path ordering* \succ_{lpo} on $\mathsf{T}_{\Sigma}(X)$ induced by \succ is defined by: $s \succ_{\text{lpo}} t$ iff

- (1) $t \in \text{var}(s)$ and $t \neq s$, or
- (2) $s = f(s_1, \dots, s_m)$, $t = g(t_1, \dots, t_n)$, and
 - (a) $s_i \succeq_{\text{lpo}} t$ for some i , or
 - (b) $f \succ g$ and $s \succ_{\text{lpo}} t_j$ for all j , or
 - (c) $f = g$, $s \succ_{\text{lpo}} t_j$ for all j , and $(s_1, \dots, s_m) (\succ_{\text{lpo}})_{\text{lex}} (t_1, \dots, t_n)$.

There are several possibilities to compare subterms in (2)(c):

compare list of subterms lexicographically left-to-right (“*lexicographic path ordering (lpo)*”, Kamin and Lévy)

compare list of subterms lexicographically right-to-left (or according to some permutation π)

compare multiset of subterms using the multiset extension (“*multiset path ordering (mpo)*”, Dershowitz)

to each function symbol f with $\text{arity}(f) \geq 1$ associate a status $\in \{\text{mul}\} \cup \{\text{lex}_{\pi} \mid \pi : \{1, \dots, n\} \rightarrow \{1, \dots, n\}\}$ and compare according to that status (“*recursive path ordering (rpo) with status*”)

The Knuth-Bendix Ordering

Let $\Sigma = (\Omega, \Pi)$ be a finite signature, let \succ be a strict partial ordering (“precedence”) on Ω , let $w : \Omega \cup X \rightarrow \mathbb{R}_0^+$ be a *weight function*, such that the following admissibility conditions are satisfied:

$$w(x) = w_0 \in \mathbb{R}^+ \text{ for all variables } x \in X; w(c) \geq w_0 \text{ for all constants } c \in \Omega.$$

If $w(f) = 0$ for some $f \in \Omega$ with $\text{arity}(f) = 1$, then $f \succeq g$ for all $g \in \Omega$.

The weight function w can be extended to terms as follows:

$$w(t) = \sum_{x \in \text{var}(t)} w(x) \cdot \#(x, t) + \sum_{f \in \Omega} w(f) \cdot \#(f, t).$$

The *Knuth-Bendix ordering* \succ_{kbo} on $T_{\Sigma}(X)$ induced by \succ and w is defined by: $s \succ_{\text{kbo}} t$ iff

- (1) $\#(x, s) \geq \#(x, t)$ for all variables x and $w(s) > w(t)$, or
- (2) $\#(x, s) \geq \#(x, t)$ for all variables x , $w(s) = w(t)$, and
 - (a) $t = x$, $s = f^n(x)$ for some $n \geq 1$, or
 - (b) $s = f(s_1, \dots, s_m)$, $t = g(t_1, \dots, t_n)$, and $f \succ g$, or
 - (c) $s = f(s_1, \dots, s_m)$, $t = f(t_1, \dots, t_m)$, and $(s_1, \dots, s_m) (\succ_{\text{kbo}})_{\text{lex}} (t_1, \dots, t_m)$.

Theorem 4.30 *The Knuth-Bendix ordering induced by \succ and w is a simplification ordering on $T_{\Sigma}(X)$.*

Proof. Baader and Nipkow, pages 125–129. □