

Universität des Saarlandes FR Informatik



Marek Kosta Christoph Weidenbach June 18, 2012

Tutorials for "Automated Reasoning" Exercise sheet 10

Exercise 10.1: (4 P)

Let a, b, c be constants, f binary function symbol and P and Q binary predicates. Apply the resolution calculus on the following two clauses:

$$C_1 = \neg Q(f(x,z), f(y,z)) \lor Q(x,y) \lor Q(z,c) \lor \neg P(x,y)$$

$$C_2 = \neg Q(f(a,c),b) \lor \neg P(a,x) \lor Q(a,b).$$

How many different possibilities to apply the resolution calculus are there? Show all resulting clauses with respective substitutions.

Exercise 10.2: (2 P)

Let $\Sigma = \{a/0, b/0, f/1, g/1; P/1, Q/1\}$ be a signature (P, Q are predicates, the other symbols)are function symbols). Find some Knuth-Bendix ordering (i.e. define weight function and precedence) in such a way that the following will hold: $Q(f(b)) \vee Q(g(b)) \succ_{\text{kbo}} P(f(a)) \vee$ $P(f(a)) \vee \neg P(g(b)) \succ_{\text{kbo}} P(g(b)) \vee P(a) \succ_{\text{kbo}} P(g(b)) \vee Q(b) \succ_{\text{kbo}} \neg P(f(a)) \vee Q(b) \succ_{\text{kbo}} \neg P(a).$ Note that for the ordering we consider the usual lifting of \succ_{kbo} to literals (negative larger) and clauses (multiset extension). Justify your definitions.

Exercise 10.3: (3 P)

Let w be a weight function defined as follows on function symbols: w(f) = 3, w(g) = 2, w(a) = 1. On predicate symbols the function w is defined as w(P) = 5, w(Q) = 4. Weight of every variable is 1. Let \succ be a precedence on symbols defined like: $P \succ Q \succ f \succ g \succ a$. Which literals in the following clauses are maximal when we consider Knuth-Bendix ordering induced by the weight function w and the precedence \succ ? Mark all the maximal literals and justify your choice.

- 1. $\neg P(g(a)) \lor P(f(f(a)))$
- 2. $P(f(x)) \lor P(g(y))$

3. $\neg P(a) \lor \neg P(f(a)) \lor Q(f(a), f(f(a)))$ 4. $\neg Q(f(x), y) \lor Q(f(f(x)), y) \lor P(x)$ 5. $\neg P(x) \lor P(f(x))$ 6. $Q(f(z)) \lor \neg P(f(g(x))) \lor \neg Q(g(f(f(y))))$

Exercise 10.4: (2 P)

Show some selection function sel such that the following set of clauses N is saturated with respect to the calculus Sup_{sel}^{\succ} for some ordering \succ . $N = \{\neg P(x) \lor Q(f(a)) \lor P(f(x)), P(g(x)) \lor P(g(f(x))) \lor \neg P(a), P(g(x)) \lor Q(y) \lor P(f(f(f(y)))) \lor \neg Q(f(y)), P(g(y)) \lor \neg Q(f(a)), Q(f(z)) \lor P(f(g(f(z))))) \lor \neg Q(f(g(f(z))))\}$. Justify your definition of the selection function.

Exercise 10.5: (*3 P*) Prove Lemma 3.28.

Exercise 10.6: (3 Bonus Points)

Consider a signature Σ . Let N be some set of ground clauses. Prove that for any ground clause C which is not a tautology (i.e. C does not contain complementary literals) over the signature Σ , $N \models C$ if and only if $N \vdash_{Res} D$ for some clause D such that $D \subseteq C$.

Submit your solution in lecture hall 001 during the lecture on June 25. Please write your name and the date of your tutorial group on your solution.

Note: Joint solutions are not permitted (work in groups is encouraged).