



Uwe Waldmann

December 1, 2011

Tutorials for “Automated Reasoning II”
Exercise sheet 5

Exercise 5.1:

Demonstrate that DPLL(T) becomes unsound if the DPLL procedure uses the pure literal rule.

Exercise 5.2:

In many applications of DPLL or DPLL(T), one does not only want a yes/no answer, but also an explanation for it. In the case of an unsatisfiable input, this explanation is an “unsatisfiable core”, i.e., a (small) subset of the input clauses that is already sufficient to show \mathcal{T} -inconsistency. How can we get an unsatisfiable core from a DPLL(T) proof?

Exercise 5.3:

Many decision procedures detect the unsatisfiability of a set of literals by iteratively deriving new literals from given literals; if an inconsistent literal is derived at the end, the input is unsatisfiable. Examples include Gaussian elimination or the Fourier-Motzkin procedure.

For such decision procedures, it is easy to generate explanations for unsatisfiability. We associate a set $E(L)$ of input literals to each literal L : for input literals L , $E(L) := \{L\}$; for literals L derived from ancestor literals L_1, \dots, L_n , $E(L) := E(L_1) \cup \dots \cup E(L_n)$. When an inconsistent literal L_0 is derived at the end, $E(L_0)$ yields the explanation.

However, the explanations computed in this way are not always minimal. Consider the following set of equations in linear rational arithmetic:

$$\begin{aligned}x - 2z &= 1 & (1) \\-x + y - 3w &= 3 & (2) \\z - 2w &= 0 & (3) \\2x - 2y + 3z &= 5 & (4)\end{aligned}$$

If we use equation (1) to eliminate x from the other equations, then (2) to eliminate y , then (3) to eliminate z , equation (4) is turned into $0 = 11$. All four equations were involved in this derivation; still $\{(1), (2), (3), (4)\}$ is *not* a minimal explanation for the contradiction. How could one efficiently find a smaller explanation? (Hint: Think about linear combinations.)

Bring your solution (or solution attempt) to the tutorial on Dec. 12.