

Automated Reasoning I*

Uwe Waldmann

Winter Term 2017/2018

Topics of the Course

Preliminaries

- abstract reduction systems
- well-founded orderings

Propositional logic

- syntax, semantics
- calculi: CDCL-procedure, ...
- implementation: 2-watched literals, clause learning

First-order predicate logic

- syntax, semantics, model theory, ...
- calculi: resolution, tableaux, ...
- implementation: sharing, indexing

First-order predicate logic with equality

- term rewriting systems
- calculi: Knuth-Bendix completion, dependency pairs

Emphasis on:

- logics and their properties,
- proof systems for these logics and their properties:
- soundness, completeness, complexity, implementation.

*This document contains the text of the lecture slides (almost verbatim) plus some additional information, mostly proofs of theorems that are presented on the blackboard during the course. It is not a full script and does not contain the examples and additional explanations given during the lecture. Moreover it should not be taken as an example how to write a research paper – neither stylistically nor typographically.

1 Preliminaries

Before we start with the main subjects of the lecture, we repeat some prerequisites from mathematics and computer science and introduce some tools that we will need throughout the lecture.

1.1 Mathematical Prerequisites

$\mathbb{N} = \{0, 1, 2, \dots\}$ is the set of natural numbers (including 0).

\mathbb{Z} , \mathbb{Q} , \mathbb{R} denote the integers, rational numbers and the real numbers, respectively.

Relations

An n -ary relation R over some set M is a subset of M^n : $R \subseteq M^n$.

For two n -ary relations R, Q over some set M , their union (\cup) or intersection (\cap) is again an n -ary relation, where

$$R \cup Q := \{ (m_1, \dots, m_n) \in M^n \mid (m_1, \dots, m_n) \in R \text{ or } (m_1, \dots, m_n) \in Q \}$$

$$R \cap Q := \{ (m_1, \dots, m_n) \in M^n \mid (m_1, \dots, m_n) \in R \text{ and } (m_1, \dots, m_n) \in Q \}.$$

A relation Q is a *subrelation* of a relation R if $Q \subseteq R$.

We often use predicate notation for relations:

Instead of $(m_1, \dots, m_n) \in R$ we write $R(m_1, \dots, m_n)$, and say that $R(m_1, \dots, m_n)$ holds or is true.

For binary relations, we often use infix notation, so

$$(m, m') \in < \Leftrightarrow <(m, m') \Leftrightarrow m < m'.$$

Words

Given a non-empty alphabet Σ , the set Σ^* of *finite words* over Σ is defined inductively by

- (i) the empty word ε is in Σ^* ,
- (ii) if $u \in \Sigma^*$ and $a \in \Sigma$ then ua is in Σ^* .

The set of *non-empty finite words* Σ^+ is $\Sigma^* \setminus \{\varepsilon\}$.

The *concatenation* of two words $u, v \in \Sigma^*$ is denoted by uv .

The length $|u|$ of a word $u \in \Sigma^*$ is defined by

- (i) $|\varepsilon| := 0$,
- (ii) $|ua| := |u| + 1$ for any $u \in \Sigma^*$ and $a \in \Sigma$.

1.2 Abstract Reduction Systems

Literature: Franz Baader and Tobias Nipkow: *Term rewriting and all that*, Cambridge Univ. Press, 1998, Chapter 2.

Throughout the lecture, we will have to work with reduction systems,

- on the object level, in particular in the section on equality,
- and on the meta level, i. e., to describe deduction calculi.

An *abstract reduction system* is a pair (A, \rightarrow) , where

- A is a non-empty set,
- $\rightarrow \subseteq A \times A$ is a binary relation on A .

The relation \rightarrow is usually written in infix notation, i. e., $a \rightarrow b$ instead of $(a, b) \in \rightarrow$.

Let $\rightarrow' \subseteq A \times B$ and $\rightarrow'' \subseteq B \times C$ be two binary relations. Then the *composition of \rightarrow' and \rightarrow''* is the binary relation $(\rightarrow' \circ \rightarrow'') \subseteq A \times C$ defined by

$$a (\rightarrow' \circ \rightarrow'') c \quad \text{if and only if} \quad a \rightarrow' b \text{ and } b \rightarrow'' c \text{ for some } b \in B.$$

\rightarrow^0	$= \{ (a, a) \mid a \in A \}$	<i>identity</i>
\rightarrow^{i+1}	$= \rightarrow^i \circ \rightarrow$	<i>$i + 1$-fold composition</i>
\rightarrow^+	$= \bigcup_{i>0} \rightarrow^i$	<i>transitive closure</i>
\rightarrow^*	$= \bigcup_{i \geq 0} \rightarrow^i = \rightarrow^+ \cup \rightarrow^0$	<i>reflexive transitive closure</i>
$\rightarrow^=$	$= \rightarrow \cup \rightarrow^0$	<i>reflexive closure</i>
\leftarrow	$= \rightarrow^{-1} = \{ (b, c) \mid c \rightarrow b \}$	<i>inverse</i>
\leftrightarrow	$= \rightarrow \cup \leftarrow$	<i>symmetric closure</i>
\leftrightarrow^+	$= (\leftrightarrow)^+$	<i>transitive symmetric closure</i>
\leftrightarrow^*	$= (\leftrightarrow)^*$	<i>refl. trans. symmetric closure or equivalence closure</i>

$b \in A$ is *reducible*, if there is a c such that $b \rightarrow c$.

b is *in normal form (irreducible)*, if it is not reducible.

c is a *normal form of b* , if $b \rightarrow^* c$ and c is in normal form.

Notation: $c = b \downarrow$ (if the normal form of b is unique).

A relation \rightarrow is called

terminating, if there is no infinite descending chain $b_0 \rightarrow b_1 \rightarrow b_2 \rightarrow \dots$

normalizing, if every $b \in A$ has a normal form.

Lemma 1.1 *If \rightarrow is terminating, then it is normalizing.*

Note: The reverse implication does not hold.

1.3 Orderings

Important properties of binary relations:

Let $M \neq \emptyset$. A binary relation $R \subseteq M \times M$ is called

reflexive, if $R(x, x)$ for all $x \in M$,

irreflexivity, if $\neg R(x, x)$ for all $x \in M$,

antisymmetric, if $R(x, y)$ and $R(y, x)$ imply $x = y$ for all $x, y \in M$,

transitive, if $R(x, y)$ and $R(y, z)$ imply $R(x, z)$ for all $x, y, z \in M$,

total, if $R(x, y)$ or $R(y, x)$ or $x = y$ for all $x, y \in M$.

A *strict partial ordering* \succ on a set $M \neq \emptyset$ is a transitive and irreflexive binary relation on M .

Notation:

\prec for the inverse relation \succ^{-1}

\succeq for the reflexive closure ($\succ \cup =$) of \succ

An $a \in M$ is called *minimal*, if there is no b in M with $a \succ b$.

An $a \in M$ is called *smallest*, if $b \succ a$ for all $b \in M \setminus \{a\}$.

Analogously:

An $a \in M$ is called *maximal*, if there is no b in M with $a \prec b$.

An $a \in M$ is called *largest*, if $b \prec a$ for all $b \in M \setminus \{a\}$.

Notation:

$M^{\prec x} = \{y \in M \mid y \prec x\}$,

$M^{\preceq x} = \{y \in M \mid y \preceq x\}$.

A subset $M' \subseteq M$ is called *downward closed*, if $x \in M'$ and $x \succ y$ implies $y \in M'$.

Well-Foundedness

Termination of reduction systems is strongly related to the concept of well-founded orderings.

A strict partial ordering \succ on M is called *well-founded* (or *Noetherian*), if there is no infinite descending chain $a_0 \succ a_1 \succ a_2 \succ \dots$ with $a_i \in M$.

Well-Foundedness and Termination

Lemma 1.2 *If $>$ is a well-founded partial ordering and $\rightarrow \subseteq >$, then \rightarrow is terminating.*

Lemma 1.3 *If \rightarrow is a terminating binary relation over A , then \rightarrow^+ is a well-founded partial ordering.*

Proof. Transitivity of \rightarrow^+ is obvious; irreflexivity and well-foundedness follow from termination of \rightarrow . \square

Well-Founded Orderings: Examples

Natural numbers. $(\mathbb{N}, >)$

Lexicographic orderings. Let $(M_1, \succ_1), (M_2, \succ_2)$ be well-founded orderings. Then let their *lexicographic combination*

$$\succ = (\succ_1, \succ_2)_{lex}$$

on $M_1 \times M_2$ be defined as

$$(a_1, a_2) \succ (b_1, b_2) \quad :\Leftrightarrow \quad a_1 \succ_1 b_1 \text{ or } (a_1 = b_1 \text{ and } a_2 \succ_2 b_2)$$

(analogously for more than two orderings)

This again yields a well-founded ordering (proof below).

Length-based ordering on words. For alphabets Σ with a well-founded ordering $>_\Sigma$, the relation \succ defined as

$$w \succ w' \quad :\Leftrightarrow \quad |w| > |w'| \text{ or } (|w| = |w'| \text{ and } w >_{\Sigma, lex} w')$$

is a well-founded ordering on the set Σ^* of finite words over the alphabet Σ (Exercise).

Counterexamples:

$(\mathbb{Z}, >)$

$(\mathbb{N}, <)$

the lexicographic ordering on Σ^*

Basic Properties of Well-Founded Orderings

Lemma 1.4 (M, \succ) is well-founded if and only if every non-empty $M' \subseteq M$ has a minimal element.

Proof. (i) “ \Leftarrow ”: Suppose that (M, \succ) is not well-founded. Then there is an infinite descending chain $a_0 \succ a_1 \succ a_2 \succ \dots$ with $a_i \in M$. Consequently, the subset $M' = \{a_i \mid i \in \mathbb{N}\}$, does not have a minimal element.

(ii) “ \Rightarrow ”: Suppose that the non-empty subset $M' \subseteq M$ does not have a minimal element. Choose $a_0 \in M'$ arbitrarily. Since for every $a_i \in M'$ there is a smaller $a_{i+1} \in M'$ (otherwise a_i would be minimal in M'), there is an infinite descending chain $a_0 \succ a_1 \succ a_2 \succ \dots$ \square

Lemma 1.5 (M_1, \succ_1) and (M_2, \succ_2) are well-founded if and only if $(M_1 \times M_2, \succ)$ with $\succ = (\succ_1, \succ_2)_{lex}$ is well-founded.

Proof. (i) “ \Rightarrow ”: Suppose $(M_1 \times M_2, \succ)$ is not well-founded. Then there is an infinite sequence $(a_0, b_0) \succ (a_1, b_1) \succ (a_2, b_2) \succ \dots$

Let $A = \{a_i \mid i \geq 0\} \subseteq M_1$. Since (M_1, \succ_1) is well-founded, A has a minimal element a_n . But then $B = \{b_i \mid i \geq n\} \subseteq M_2$ can not have a minimal element, contradicting the well-foundedness of (M_2, \succ_2) .

(ii) “ \Leftarrow ”: obvious. \square

Monotone Mappings

Let (M_1, \succ_1) and (M_2, \succ_2) be strict partial orderings. A mapping $\varphi : M_1 \rightarrow M_2$ is called *monotone*, if $a \succ_1 b$ implies $\varphi(a) \succ_2 \varphi(b)$ for all $a, b \in M_1$.

Lemma 1.6 If φ is a monotone mapping from (M_1, \succ_1) to (M_2, \succ_2) and (M_2, \succ_2) is well-founded, then (M_1, \succ_1) is well-founded.

Well-founded Induction

Theorem 1.7 (Well-founded (or Noetherian) Induction) *Let (M, \succ) be a well-founded ordering, let Q be a property of elements of M .*

If for all $m \in M$ the implication

*if $Q(m')$ for all $m' \in M$ such that $m \succ m'$,¹
then $Q(m)$.²*

is satisfied, then the property $Q(m)$ holds for all $m \in M$.

Proof. Let $X = \{m \in M \mid Q(m) \text{ false}\}$. Suppose, $X \neq \emptyset$. Since (M, \succ) is well-founded, X has a minimal element m_1 . Hence for all $m' \in M$ with $m' \prec m_1$ the property $Q(m')$ holds. On the other hand, the implication which is presupposed for this theorem holds in particular also for m_1 , hence $Q(m_1)$ must be true so that m_1 can not be in X . *Contradiction.* \square

Well-founded Recursion

Let M and S be sets, let $N \subseteq M$, and let $f : M \rightarrow S$ be a function. Then the *restriction* of f to N , denoted by $f|_N$, is a function from N to S with $f|_N(x) = f(x)$ for all $x \in N$.

Theorem 1.8 (Well-founded (or Noetherian) Recursion) *Let (M, \succ) be a well-founded ordering, let S be a set. Let ϕ be a binary function that takes two arguments x and g and maps them to an element of S , where $x \in M$ and g is a function from $M^{\prec x}$ to S .*

Then there exists exactly one function $f : M \rightarrow S$ such that for all $x \in M$

$$f(x) = \phi(x, f|_{M^{\prec x}})$$

Proof. The proof consists of four parts.

Part 1: For every downward closed subset $N \subseteq M$ there is *at most one* function $f : N \rightarrow S$ such that $f(x) = \phi(x, f|_{M^{\prec x}})$.

Proof: Assume that there exist a downward closed subset $N \subseteq M$ and two *different* functions f_1 and f_2 from N to S with this property. Therefore, the set $N' := \{x \in N \mid f_1(x) \neq f_2(x)\}$ is non-empty. By well-foundedness, N' has a minimal element y . By minimality of y , $f_1|_{M^{\prec y}} = f_2|_{M^{\prec y}}$. Therefore $f_1(y) = \phi(y, f_1|_{M^{\prec y}}) = \phi(y, f_2|_{M^{\prec y}}) = f_2(y)$, contradicting the assumption.

¹induction hypothesis

²induction step

Part 2: If N_1 and N_2 are downward closed subsets of M and the functions $f_1 : N_1 \rightarrow S$ and $f_2 : N_2 \rightarrow S$ satisfy $f_i(x) = \phi(x, f|_{M^{\prec x}})$ for all $x \in N_i$ ($i = 1, 2$), then $f_1(x) = f_2(x)$ for all $x \in N_1 \cap N_2$.

Proof: Define $N_0 := N_1 \cap N_2$ and $f'_i = f_i|_{N_0}$ for $i = 1, 2$. Clearly N_0 is downward closed and for all $x \in N_0$ and $i = 1, 2$ we have $f'_i(x) = f_i(x) = \phi(x, f_i|_{M^{\prec x}}) = \phi(x, f'_i|_{M^{\prec x}})$. By part 1, there is at most one function from N_0 to S with this property, so $f'_1 = f'_2$, and therefore $f_1(x) = f_2(x)$ for all $x \in N_1 \cap N_2$.

Part 3: For every $y \in M$ there exists a function $f_y : M^{\preceq y} \rightarrow S$ such that $f_y(x) = \phi(x, f_y|_{M^{\prec x}})$ for all $x \in M^{\preceq y}$.

Proof: We use well-founded induction over \succ . Let $y \in M$. By the induction hypothesis, for every $z \prec y$ there exists a function $f_z : M^{\preceq z} \rightarrow S$ such that $f(x) = \phi(x, f|_{M^{\prec x}})$ for all $x \in M^{\preceq z}$. By part 2, all functions f_z agree on the intersections of their domains. Define the function $f_y : M^{\preceq y} \rightarrow S$ by $f_y(x) = f_x(x)$ for $x \prec y$ and by $f_y(y) = \phi(y, f_y|_{M^{\prec y}})$.

Part 4: There exists a function $f : M \rightarrow S$ such that $f(x) = \phi(x, f|_{M^{\prec x}})$ for all $x \in M$.

Proof: Define $f : M \rightarrow S$ by $f(x) = f_x(x)$.

The claim of the theorem follows now from part 1 (for $N := M$) and part 4. □

The well-founded recursion scheme generalizes terminating recursive programs.

Note that functions defined by well-founded recursion need *not* be computable, in particular since for many well-founded orderings the sets $M^{\prec x}$ may be infinite.

1.4 Multisets

Let M be a set. A *multiset* S over M is a mapping $S : M \rightarrow \mathbb{N}$. We interpret $S(m)$ as the number of occurrences of elements m of the base set M within the multiset S .

Example. $S = \{a, a, a, b, b\}$ is a multiset over $\{a, b, c\}$, where $S(a) = 3$, $S(b) = 2$, $S(c) = 0$.

We say that m is an *element* of S , if $S(m) > 0$.

We use set notation (\in , \subseteq , \cup , \cap , etc.) with analogous meaning also for multisets, e. g.,

$$\begin{aligned}
 m \in S & \quad :\Leftrightarrow \quad S(m) > 0 \\
 (S_1 \cup S_2)(m) & \quad := \quad S_1(m) + S_2(m) \\
 (S_1 \cap S_2)(m) & \quad := \quad \min\{S_1(m), S_2(m)\} \\
 (S_1 - S_2)(m) & \quad := \quad \begin{cases} S_1(m) - S_2(m) & \text{if } S_1(m) \geq S_2(m) \\ 0 & \text{otherwise} \end{cases} \\
 S_1 \subseteq S_2 & \quad :\Leftrightarrow \quad S_1(m) \leq S_2(m) \text{ for all } m \in M
 \end{aligned}$$